

Diving Eggs


A Demonstration of Inertia

Introduction

Students experience the effects of inertia every day—riding in a car, playing sports, even when picking up their backpacks. Present a more dramatic demonstration of Newton’s first law by safely dropping three raw eggs into glasses of water, without touching the eggs! Applause is guaranteed!

Concepts

- Balanced forces
- Inertia
- Newton’s first law of motion



Materials

- | | |
|---|---------------------------------------|
| Eggs, raw, 3 | Drinking glasses, 3 |
| Water, tap | Pie pan or pizza pan, sturdy aluminum |
| Broom | Table |
| Cardboard tubes, 3 (empty toilet paper tubes) | Towel |
| Demonstration tray or jelly roll pan | |

Safety Precautions

Take care when handling raw eggs. Wash hands thoroughly with soap and water after handling eggs. Clean up spills immediately. Wear safety glasses. Observers should be a safe distance away from the trajectory of the pie pan. Follow all laboratory safety guidelines.

Procedure

1. Obtain three identical tall drinking glasses, large enough for an egg to fit in sideways.
2. Fill each glass about three-fourths full with tap water.
3. Place a demonstration tray or jelly roll pan 1 cm from the edge of a table. *Note:* The bottom of the tray must be completely flat.
4. Place two of the glasses in the tray next to the edge. The rim of the glasses should be near, but not extending over, the edge of the table.
5. Place the third glass between the other two, forming a triangle shape. See Figure 1.
6. Obtain a sturdy aluminum pie plate or a small aluminum round pizza pan.
7. Center the aluminum pan on top of the three glasses. The edge of the pan should extend slightly beyond the edge of the table.
8. Obtain three empty cardboard toilet paper tubes.
9. Stand the tubes vertically on the aluminum pan, centering one tube over each glass (see Figure 2).
10. Obtain three raw eggs. *Note:* For practice, hard boiled eggs may be used. Once the demonstration has been perfected, use raw eggs.
11. Balance one egg on top of each cardboard tube, laying the egg lengthwise across the top of the tube. Do not fit the narrow end of the egg into the tube; the egg should just rest on top (see Figure 2).
12. Obtain a broom and place the bristles on the floor by the table with the handle extending upward.
13. Holding on to the broom handle, and facing the egg setup on top of the table, step on the bristles of the broom with one foot.




Figure 1. Overhead view




Figure 2.

14. Pull back on the broom handle like a lever. See Figure 3.
15. *Without letting go of the broom handle*, carefully test to see if the handle will hit the aluminum pan *between* the two closer glasses when the handle is released. If not, adjust the position of the broom accordingly. The handle should not hit any of the glasses or the tray.
16. Continuing to step on the bristles, pull the broom handle back.
17. Let go of the broom handle and watch as the eggs “dive” safely into the glasses of water.



Figure 3.

Disposal

Please consult your current *Flinn Scientific Catalog/Reference Manual* for general guidelines and specific procedures governing the disposal of laboratory wastes. Eggs may be disposed of in the trash according to Flinn Suggested Disposal Method #26a. Egg whites and yolks from broken eggs may be disposed of down the drain with plenty of water according to Flinn Suggested Disposal Method #26b. If necessary, sterilize the glasses and demonstration tray with a 10% bleach solution, then rinse with water.

NGSS Alignment

This laboratory activity relates to the following Next Generation Science Standards (2013):

Disciplinary Core Ideas: Middle School

- MS-PS2 Motion and Stability: Forces and Interactions
 - PS2.A: Forces and Motion
 - PS2.B: Types of Interactions

Disciplinary Core Ideas: High School

- HS-PS2 Motion and Stability: Forces and Interactions
 - PS2.A: Forces and Motion
 - PS2.B: Types of Interactions

Science and Engineering Practices

- Developing and using models
- Planning and carrying out investigations

Crosscutting Concepts

- Patterns
- Cause and effect

Tips

- Practice with other objects before using raw eggs. Hard boiled eggs, rubber balls, or any other objects similar in size and mass may be used.
- Thin aluminum pie pans may dent with the force of the broom handle, resulting in a less elastic collision. These pans are not recommended for this demonstration.
- Diving Eggs is available from Flinn Scientific as a Super Value Kit, “Diving Eggs Inertia Challenge” (Catalog No. AP7419). The kit includes weighted plastic eggs for practice.
- Using clear drinking glasses allows students to see the eggs after they have dropped into the water. Large beakers may be used instead of drinking glasses.
- Videotape the demonstration, and then play it back in slow motion.

Discussion

Newton’s first law of motion states that an object at rest tends to stay at rest unless a net force acts on it. This law is also known as the law of inertia. *Inertia* is the tendency of an object to resist change in motion. Inertia is directly related to mass—the greater the mass of an object, the greater its inertia. In the “Diving Eggs” setup, all the forces are balanced with all objects at rest. A net horizontal force is supplied by the moving broom handle. This force acts upon the aluminum pan, which then accelerates in the direction of the applied force. When the edge of the pan hits the cardboard tubes, the tubes accelerate in the direction of the force also. The eggs do not move with the tubes because of their greater inertia. Since no horizontal force acts on the eggs, however, once the force holding them up is gone, the only remaining force is gravity, and the eggs drop into the glasses of water.

Materials for *Diving Eggs* are available from Flinn Scientific, Inc.

| Catalog No. | Description |
|-------------|---|
| AP7419 | Diving Eggs Inertia Challenge—Super Value Kit |
| GP1025 | Beaker, 400-mL |

Consult your *Flinn Scientific Catalog/Reference Manual* for current prices.